

A symmetry problem ^{*†}

A.G. Ramm

Mathematics Department, Kansas State University,
Manhattan, KS 66506-2602, USA

ramm@math.ksu.edu

Abstract

The following result is proved:

Theorem. Let $D \subset \mathbb{R}^3$ be a bounded domain homeomorphic to a ball, $|D|$ be its volume, $|S|$ be the surface area of its smooth boundary S , $D \subset B_R := \{x : |x| \leq R\}$, and H_R is the set of all harmonic in B_R functions. If

$$\frac{1}{|D|} \int_D h dx = \frac{1}{|S|} \int_S h ds \quad \forall h \in H_R,$$

then D is a ball.

1 Introduction

Consider a bounded domain $D \subset \mathbb{R}^n$, $n = 3$, with a boundary S . The exterior domain is $D' = \mathbb{R}^3 \setminus D$. Assume that S is Lipschitz. Let N be the exterior unit normal to S . Our result is:

Theorem. *Let D be a bounded domain homeomorphic to a ball, $|D|$ be its volume, $|S|$ be the surface area of its smooth boundary S , $D \subset B_R := \{x : |x| \leq R\}$, and H_R is the set of all harmonic in B_R functions. If*

$$\frac{1}{|D|} \int_D h dx = \frac{1}{|S|} \int_S h ds \quad \forall h \in H_R, \tag{1}$$

then D is a ball.

We prove this theorem using the result from [1] and the following lemmas:

Lemma 1. *The set $\{h|_S\}_{\forall h \in H_R}$ is dense in $L^2(S)$.*

For convenience of the reader let us formulate the result from [1]:

^{*}key words: symmetry for PDE, harmonic functions

[†]AMS subject classification: 35R30

Lemma 2. *If D is homeomorphic to a ball and*

$$\Delta u = \frac{1}{|D|} \text{ in } D, \quad u|_S = 0, \quad u_N|_S = \text{const},$$

then D is a ball.

The result is valid for $D \in \mathbb{R}^n$, $n \geq 2$, and its proof is essentially the same. If D is a ball, then formula (1) holds and both integrals are equal to $h(0)$, where the origin is at the center of the ball (mean-value theorem for harmonic functions).

In the next Section proofs are given.

2 Proofs

Proof of Lemma 1. Assume the contrary. Then there exists an f such that $\int_S f h ds = 0$ for all $h \in H_R$. Take $h = \frac{1}{4\pi|x-y|}$ where $y \in B'_R := \mathbb{R}^3 \setminus B_R$ is arbitrary. Then

$$w(y) := \int_S \frac{f ds}{4\pi|s-y|} = 0 \quad \forall y \in B'_R.$$

Since w is harmonic in D' , it follows that $w = 0$ in D' . Since w is harmonic in D , it follows that $w = 0$ in D . By the jump formula for the normal derivative of the simple-layer potential, one concludes that $f = 0$. Lemma 1 is proved. $\square$

Proof of Theorem 1. Let u be the unique solution to the problem:

$$\Delta u = \frac{1}{|D|} \quad \text{in } D, \quad u|_S = 0. \quad (2)$$

Multiply (2) by an arbitrary $h \in H_R$, integrate over D , then by parts, use the boundary condition (2) and get:

$$\frac{1}{|D|} \int_D h dx = \int_S u_N h ds \quad \forall h \in H_R. \quad (3)$$

By (1) this implies:

$$\int_S (u_N - \frac{1}{|S|}) h ds = 0 \quad \forall h \in H_R. \quad (4)$$

By Lemma 1, equation (4) implies

$$u_N = \frac{1}{|S|} \quad \text{on } S. \quad (5)$$

By Lemma 2, equations (2) and (5) imply that D is a ball. Theorem 1 is proved. $\square$

References

- [1] J. Serrin, A symmetry problem in potential theory, Archive for rational mechanics and analysis, 43, (1971), 304 - 318.